

Recombinant Mouse EFNA1 (C-Fc-6His)

Catalog #	EPT280	
Expression Host	Human Cells	
DESCRIPTION	Recombinant Mouse Ephrin-A1 is produced by our	
	Mammalian expression system and the target gene	
	encoding Asp19-Ser182 is expressed with a Fc, 6His	
	tag at the C-terminus.	
Accession	P52793	
Synonyms	EPH-related receptor tyrosine kinase ligand 1;	
	Immediate early response protein B61;Epgl1; Epl1;	
	Lerk1	
Mol Mass	47.3 KDa	
AP Mol Mass	45-65 KDa, reducing conditions	
Purity	Greater than 90% as determined by reducing	
	SDS-PAGE.	
Endotoxin	Less than 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL	
	test.	
FORMULATION	Lyophilized from a 0.2 µm filtered solution of PBS, pH	



+86-27-59760950

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C

www.elkbiotech.com

ELKbio@ELKbiotech.com



RECONSTITUTION Always centrifuge tubes before opening.Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. SHIPPING The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. Lyophilized protein should be stored at < -20 °C, STORAGE though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20° C for 3 months. BACKGROUND Ephrin-A1 is a cell membrane protein and contains 1 ephrin RBD (ephrin receptor-binding) domain. EFNA1 belongs to the ephrin (EPH) family. The ephrins and EPH-related receptors comprise the largest subfamily of receptor protein-tyrosine kinases and have been mediating developmental implicated in events,



+86-27-59760950

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.Q

www.elkbiotech.com

ELKbio@ELKbiotech.com



especially in the nervous system and in erythropoiesis. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. This gene encodes an EFNA class ephrin which binds to the EPHA2, EPHA4, EPHA5, EPHA6, and EPHA7 receptors. Two transcript variants that encode different isoforms were identified through sequence analysis. It belongs to the ephrin family and contains 1 ephrin RBD (ephrin receptor-binding) domain.

kDa	МК	R
120 90 60		
40		-
30	- Second	
20	Band er	
14		

SDS-PAGE



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.Q